


## Supplementary material (JCEM / jc.2019-01294)

### RESIDUAL ADRENAL FUNCTION IN AUTOIMMUNE ADDISON'S DISEASE – EFFECT OF DUAL THERAPY WITH RITUXIMAB AND DEPOT TETRACOSACTIDE


Catherine Napier<sup>1</sup>, Earn H Gan<sup>1</sup>, Anna L Mitchell<sup>1</sup>, Lorna C Gilligan<sup>2</sup>, D Aled Rees<sup>3</sup>, Carla Moran<sup>4</sup>, Krishna Chatterjee<sup>4</sup>, Bijay Vaidya<sup>5</sup>, R Andrew James<sup>1</sup>, Yaasir Mamoojee<sup>1</sup>, Simon Ashwell<sup>6</sup>, Wiebke Arlt<sup>2,7</sup>, Simon HS Pearce<sup>1</sup>

#### Supplementary figure

Panel A.



Panel B.



**Supplementary figure legend:** Panel A shows serum immunoglobulin A, G and M concentrations at baseline, 12- and 48- weeks in the 13 study participants. LLN refers to the lower limit of the reference range. Panel B shows the % of CD19+ lymphocytes out of the total lymphocyte population measured by flow cytometry during the study.

**Supplementary Table**

<b>Serious Adverse Events*</b>	<b>Number of events</b>	<b>Number of participants (%)</b>
Diarrhoea, vomiting, gastrointestinal illness	3	3 (23%)
Facial laceration (vehicle accident)	1	1 (8%)
*All of the Serious adverse events were classified as SAEs because of hospital attendance. None were classified as treatment-related by the investigators		
<b>Other Adverse Events</b> (excluding inject-site reactions)		
Back pain	84	9 (69%)
Headache		
Fatigue		
Cough		
Poor concentration		
Sinusitis		
Sore throat		
Tonsillitis		
Ear infection		
Chest infection		
Urinary tract infection		
Itchy skin		
Conjunctivitis		
Diarrhoea		
Abdominal pain		
Dizziness		
Dehydration		
Cut to hand		
Weight loss		